

Determining the effect of *clpA* in the natural transformation ability of *Aeromonas salmonicida*

Kristen Clemons and Dr. Jennifer Huddleston

Biology Department, Abilene Christian University, Abilene Texas 79699

Abstract

Members of the genus *Aeromonas* are ubiquitous in water and can be identified as Gram-negative, rod-shaped, facultatively anaerobic bacteria. These bacteria are opportunistic extraintestinal and intraintestinal pathogens that can cause disease year-round with varying severity. *Aeromonas salmonicida* is a naturally transformable species. Natural transformation is a six-part process that results in a bacterium incorporating free DNA into its genome. It is possible that aeromonads can become antibiotic resistant through this mechanism. Having been indicated as affecting the natural transformation process in the genus *Aeromonas*, the effect of the proteolytic molecular chaperone ClpA on the natural transformation ability of *Aeromonas* was studied in this investigation. This was accomplished through the creation of a *clpA* mutant fragment in which *clpA* was replaced with a gentamycin resistance cassette. This fragment was assembled using PCR and SOE-PCR. Since the mutant fragment has been constructed, the project is now focusing on the gene replacement in the wild-type *Aeromonas* with the intention of making a comparison between the wild-type and mutant strains which will enable the determination of the effect of ClpA on the natural transformation ability of *Aeromonas salmonicida*.

Introduction

Bacteria in the genus *Aeromonas* are ubiquitous in aquatic environments. These mesophilic bacteria have been found in domestic sewage, rivers receiving sewage discharges, clean rivers, lakes and storage reservoirs, as well as drinking water in distribution systems (1). Aeromonads are known to cause a variety of diseases in humans. These are pathogens that can infect healthy humans of all ages with varying degrees of severity through a number of different modes of infection (2, 3, 4). In addition, Sakai (5) and Huddleston et al (6) reported that *Aeromonas salmonicida* is capable of achieving competence and undergoing natural transformation. In this process, a bacterial cell obtains free DNA from the environment to use for survival in its environment. The cell requires the participation of certain cellular functions that are provided by genes located throughout the bacterial chromosome and that may only be expressed under particular environmental conditions (7). The competence-inducing conditions for *Aeromonas* are known to include nutrient-limiting conditions and stationary phase of growth (6) Transformation by free DNA is more likely to occur in habitats with high bacterial concentrations. This environment exists in the intestines of insects, worms, and warm-blooded animals including the human stomach and intestines (7). The presence of antibiotics is stressful enough to induce competence in bacteria. Antibiotics create an environment in which antibiotic-resistant populations are selected. These antibiotic-resistant populations then expand clonally and through horizontal gene transfer (8). Since *Aeromonas* can survive in the intestines and causes gastrointestinal infections, the prospect of this bacteria becoming antibiotic resistant increases, and therefore, is more difficult to treat. If the processes of horizontal gene transfer were understood, the acquiring of antibiotic resistance could be averted along with the loss of human lives. Clp proteins refold, remodel and degrade proteins damaged under stress conditions (9, 10). It is reasonable to conclude that there could be an increase in the presence or activity of ClpA at the time of natural transformation. To test for the importance of ClpA in the natural transformation ability of *Aeromonas salmonicida*, a mutant fragment containing a gentamycin resistance cassette was constructed to replace *clpA* in the wild-type *Aeromonas* genome.




Figure 1. Gene Map of the *clpA* region in the *Aeromonas* wild type genome.

Materials and Methods

Mutant Fragment Construction. Using the previously extracted DNA from *Aeromonas salmonicida* strain #92, approximately 500 base pairs from the downstream and upstream *clpA* region were amplified through Polymerase Chain Reaction (PCR) using the designed primers. The gentamycin resistance cassette (Gm^R) was amplified through PCR from *Aeromonas* strain #771 which has been engineered to contain the gene for gentamycin resistance (Gm^R). The PCR products were purified using the Zymo Research DNA Clean & Concentrator™-25 kit (Irvine, CA). Splicing overlap extension polymerase chain reaction (SOE-PCR) (11) was used to combine the three amplified pieces. First, the gentamycin resistance cassette (Gm^R) and the upstream *clpA* region were combined using this method. Then, the downstream *clpA* region was attached to this product using SOE-PCR followed by a gel extraction to ensure the isolation of the pure mutant fragment. This gel extraction was performed using a 50 mL 2.0% agarose gel and the Zymoclean™ Gel DNA Recovery Kit (Irvine, CA).

Figure 2. SOE-PCR Scheme: Splicing Overlap Extension Polymerase Chain Reaction




Figure constructed by K. Clemons and J. Huddleston, 2014




Figure 3. Gels depicting Mutant Fragment Construction.

Left: Gel electrophoresis of PCR amplification of *clpA* flanking regions and the Gm^R cassette.
Middle: SOE-PCR combination and amplification of upstream *clpA* region/Gm^R
Right: PCR amplification of the *clpA* mutant fragment

Results and Conclusion

As can be seen in Figure 3, the *clpA* mutant fragment was constructed. Constructing the *clpA* mutant fragment is the first step to achieving an understanding of the role of ClpA in the natural transformation ability of *Aeromonas salmonicida*. Now that this has been accomplished, gene replacement of the *Aeromonas* wild-type is possible. The gene replacement will allow for a comparison of the wild-type and mutant bacteria which will enable determination of the extent of involvement of ClpA in the natural transformation ability of *Aeromonas salmonicida*.




Figure 4. Gene Map of *clpA* mutant fragment.

Discussion

The downstream and upstream *clpA* regions of the mutant fragment act as recognition regions for the *clpA* site in the genome. The homologous regions should anneal to each other, causing the mutant fragment to replace *clpA* in the wild-type *Aeromonas* genome. The Gm^R serves as a selectable marker. The study of natural transformation in the genus *Aeromonas* is only beginning. Huddleston (12) implicated four genes in the ability of aeromonads to naturally transform, but Maughan and Redfield (13) related that 30 genes have been identified in *Haemophilus influenza* as being responsible for its natural transformation ability. The construction of this *clpA* mutant fragment has paved the way for projects to come. In addition to launching the creation of the *clpA* *Aeromonas* mutant, this method can be used to construct mutant fragments for the study of *clpS* and *clpA/clpS* in the natural transformation ability of *Aeromonas* as well as other genes suspected of involvement in its natural transformation ability.

Citations

1. Holmes P, Nicolls LM, Sartory DP. 1996. The Ecology of Mesophilic *Aeromonas* in the Aquatic Environment, p. 128-150. In Austin, B., Altwegg, M., Gosling, P.J., and Joseph, S. (Eds.), The Genus *Aeromonas*. John Wiley & Sons Ltd, West Sussex, England.
2. Janda, MJ and Abbott SL. 1996. Human Pathogens, p. 151-173. In Austin, B., Altwegg, M., Gosling, P.J., and Joseph, S. (Eds.), The Genus *Aeromonas*. John Wiley & Sons Ltd, West Sussex, England.
3. Janda JM and Abbott SL. 2010. The Genus *Aeromonas*: Taxonomy, Pathogenicity, and Infection. Clin. Microbiol. 23: 35-73.
4. Joseph, S. W. 1996. *Aeromonas* Gastrointestinal Disease: A Case Study in Causation?, p. 311-355. In Austin, B., Altwegg, M., Gosling, P.J., and Joseph, S. (Eds.), The Genus *Aeromonas*. John Wiley & Sons Ltd, West Sussex, England.
5. Sakai DK. 1987. Transformation of a Nonpathogenic *Aeromonas salmonicida* Mutant with Intracellular and Intergenic Bacterial DNA Fragments in a Laboratory Model of River Sediments. Sci. Rep. Hokkaido Fish Hatch. 53: 1141-1149.
6. Huddleston JR, Brokaw JM, Zak JC, and Jeter RM. 2013. Natural transformation as a mechanism of horizontal gene transfer among environmental *Aeromonas* species. Syst. Appl. Microbiol. 36: 224-234.
7. Lorenz MG and Wackernagel W. 1994. Bacterial gene transfer by natural genetic transformation in the environment. Microbiol. Rev. 58: 563-602.
8. Huddleston, J. R. 2014. Horizontal gene transfer in the human intestines: potential spread of antibiotic resistance genes. Infect. Drug Resist. 7: 1-10.
9. Pak M. and Wickner S. 1997. Mechanism of protein remodeling by ClpA chaperone. Proc. Natl. Acad. Sci. USA. 94: 4901-4906.
10. Wawrzynow A, Banecki B. and Zylcz M. 1996. The Clp ATPases define a novel class of molecular chaperones. Mol. Microbiol. 21: 895-899.
11. Higuchi R, Krummel B. and Saiki R. 1988. A general method of in vitro preparation and specific mutagenesis of DNA fragments: study of protein and DNA interactions. Nucleic Acids Res. 16: 7351-7367.
12. Huddleston JR. 2008. Natural Transformation in *Aeromonas* species. Ph. D. dissertation. Texas Tech University, Lubbock, TX.
13. Maughan H. and Redfield RJ. 2009. Extensive variation in natural competence in *Haemophilus influenzae*. Evolution. 63: 1852-1866.

Acknowledgements

This research was supported by funds from the McNair Scholars Program. Travel was supported by ACU Office of Undergraduate Research.